

How to use your WAINUA autoinjector

WAINUA is a once-monthly injection you can take at home or on the go (proper storage required) for the treatment of hereditary transthyretin-mediated amyloidosis (hATTR-PN) in adults.

Your healthcare provider should have shown you or your caregiver how to use the autoinjector the right way. Use this guide as a reminder. There are 3 sections of steps:

1. Prep 2. Inject 3. Check


For complete instructions, please read the "Instructions for Use" in the full Prescribing Information.

Important Information you need to know before using WAINUA

- WAINUA is for injection under the skin (subcutaneous use) only
- Each WAINUA autoinjector contains 1 dose and can only be used 1 time
- **Do not** share your autoinjector with anyone



1. Prep your supplies and injection site



If your autoinjector was in the refrigerator, **keep it in the original carton after removing it from the refrigerator and let it come to room temperature (up to 86 °F or 30 °C) for 30 minutes before injecting.**

Do not warm up in any other way (eg, in a microwave, hot water, or near other heat sources). Keep it away from light or direct sunlight.


Take your autoinjector out of its carton. **Do not use autoinjector if it has:**

- ⊗ been frozen
- ⊗ been dropped, damaged, or appears to be tampered with
- ⊗ passed the expiration date (EXP)



The liquid in the viewing window should be clear and colorless to slightly yellow. **Do not use if the liquid:**

- ⊗ looks cloudy
- ⊗ is discolored
- ⊗ contains large particles



- Self-injection sites: abdomen (stomach), front of thigh
- Only caregivers should inject into the back of the upper arm

Choose an injection site. **Do not inject:**

- ⊗ into the 2-inch (5-cm) area around your belly button
- ⊗ into skin that is red, warm, tender, bruised, scaly, or hard
- ⊗ into scars, damaged, discolored, or tattooed skin
- ⊗ through clothing

Wash your hands well with soap and water.

Clean your injection site by rubbing an alcohol wipe or with soap and water in a circular motion. Let it air-dry.

Do not touch the injection site once it is clean.

2. Inject WAINUA




Hold the autoinjector body with one hand, with the clear cap facing away from you.

Pull the cap straight off; **do not twist it off**. Throw away the cap. **Do not** recap the autoinjector. This could cause the medicine to come out too soon or damage the autoinjector.

Don't see the needle? That's normal. The needle remains hidden inside the orange needle guard.

Do not touch the needle or push the orange needle guard against your hand or finger.



Place the orange needle guard flat on your injection site at a **90-degree angle** with the viewing window facing you.



Firmly press and hold the autoinjector against your skin for 10 seconds to make sure the full dose has been given. You may hear a "click" when the injection starts. The orange plunger rod will start to move down in the viewing window.

You may hear a second "click," but the injection is not done yet—**keep holding for the full 10 seconds. Do not move or change the position of the autoinjector after the injection has started.**

3. Check that you injected all the medicine



After you have completed your injection, lift the autoinjector straight up. Check the viewing window to make sure all the medicine has been injected. If the orange plunger rod does not fill the viewing window, you may not have received the full dose. If this happens or you have other concerns, contact your healthcare provider. **Do not try to inject again.**

Immediately put the used autoinjector in an FDA-cleared sharps disposal container. **Do not throw into household trash.**

There may be a small amount of blood or liquid where you injected; this is normal. If needed, press a clean cotton ball or gauze on the area and apply a small bandage.



Watch our instructional video for step-by-step guidance on how to take WAINUA.

Please see Important Safety Information on page 4.

WAINUA is a registered trademark of the AstraZeneca group of companies.
 ©2026 AstraZeneca. All rights reserved. US-107894 Last Updated 2/26

Frequently asked questions about taking WAINUA

Q: How will I receive my autoinjector and supplies?

A: Your autoinjector and sharps container will be mailed to you by a specialty pharmacy. They might call or text to schedule your delivery. **Enroll in WAINUA WAY to receive a welcome kit with alcohol wipes and helpful tools to get you started.** Visit **WAINUAWay.com** to learn more.

Q: What if my autoinjector is damaged?

A: Do not use a damaged autoinjector. **Call 1-844-2-WAINUA (844-292-4682)** for a replacement.

Q: I don't see a needle. Is my autoinjector broken?

A: The needle is there; it's just hidden inside the orange needle guard as a safety feature. (Don't try to remove the orange safety guard.)

Q: How do I store my autoinjector?

A: Store WAINUA in a refrigerator at 36 °F to 46 °F (2 °C to 8 °C) in its original carton until you are ready to use it. **Do not** freeze. WAINUA may be kept at room temperature—up to 86 °F (30 °C)—for up to a maximum of 6 weeks, protected from light in its original carton. **Do not** let WAINUA reach temperatures above 86 °F (30 °C) or expose to heat. Protect from light. If you do not use WAINUA kept at room temperature within 6 weeks, throw it away.

Q: Can I travel with my autoinjector?

A: Always discuss travel with your healthcare provider first. You can travel with your autoinjector if you store it as described in the Instructions for Use.

Q: What are the most common side effects of WAINUA?

A: The most common side effects include decreased vitamin A and vomiting. These are not all the possible side effects of WAINUA. Talk to your healthcare provider about side effects that you experience.

Q: What if I missed my injection date?

A: If you miss a dose, take the missed dose as soon as possible. Then take WAINUA 1 month from the date of your last dose to get back on a monthly dosing schedule. If you have questions about your schedule, ask your healthcare provider.

Q: Can someone help me take WAINUA?

A: Yes, the steps are the same. Remember, they should be trained by your healthcare provider and read the Instructions for Use that came with your autoinjector first.

Caregivers can inject into the back of your upper arm, your stomach (abdomen), or the front of your thigh. Watch our instructional video at **WAINUA.com/HowToVideo** for more information.

**Have more questions? Visit [WAINUA.com](https://www.wainua.com)
or call 1-844-2-WAINUA (844-292-4682)**

IMPORTANT SAFETY INFORMATION

WAINUA may cause side effects, including:

- **Low vitamin A level**, which is a serious, but common side effect of treatment with WAINUA. Your doctor should tell you to take vitamin A supplements. **Do not** take more than the amount of vitamin A your doctor has recommended. Call your doctor if you develop eye problems such as difficulty seeing at night or in low-lit areas (night blindness), or dry eyes. If you develop eye problems, your doctor should refer you to an eye doctor

The most common side effects include decreased vitamin A and vomiting.

Before you take WAINUA, tell your healthcare provider:

- **if you are pregnant or breastfeeding, or plan to become pregnant or breastfeed.** It is not known if WAINUA can harm your baby. Changes in vitamin A levels and vitamin A supplementation related to use of WAINUA may harm your unborn baby. It is not known if WAINUA passes into breast milk
- **about all the medicines you take**, including prescription and nonprescription medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take vitamin A or beta-carotene supplements

APPROVED USES

WAINUA injection, for subcutaneous use, 45 mg is a prescription medicine used to treat adults with polyneuropathy of hereditary transthyretin-mediated amyloidosis.

Please see full [Prescribing Information](#), including [Patient Information](#), for WAINUA.

You may [report side effects related to AstraZeneca products](#) .